

JK Chrome | Employment Portal



Rated No.1 Job Application of India


Sarkari Naukri
Private Jobs
Employment News
Study Material
Notifications


















JK Chrome

jk chrome Contains ads



www.jkchrome.com | Email : contact@jkchrome.com

Classification of Plantae

The study of different types of Trees, plants is called Botany. Theophrastus is called the father of Botany.

1. Classification of Plantae

In the year 1883, Eichler has classified the Botanical world as under:-

I. Cryptogamus plants

There is no flower and seed in these type of plants. These are classified into the following groups:

Thalophyta:

- 1. This is the largest group of the plant kingdom.
- 2. The body of the plants in this group is thalus like i.e., plant are not differentiated into root, stem and leaves etc.
- There is no conducting tissue. It is divided into two groups.
 (a) Algae and (b) Fungi

(a) Algae

- 1. The study of algae is called Phycology.
- 2. The algae normally have chlorophyll and autrotrophic mode of nutrition.
- 3. Its body is thalus like. It may be unicellular, colonial or filamentous.

Useful Algae

- **1. As a food :** Porphyra, Ulva, Surgassum, Laeminaria, Nostoc etc.
- 2. In making lodine: Laeminaria, Fucus, Echlonia etc.
- 3. As a manure: Nostoc, Anabina, kelp etc.
- **4. In making medicines :** Chloreloline from Chlorella and Tincher iodine is made from Laminaria.
- **5.** In research works : Chlorella Acitabularia, Belonia etc.

Note: An astronaut can get protein food, water and oxygen by sowing the chlorella algae in the tank of the aircraft so chlorella is known as space algae.

(b) Fungi

- **1.** Study of fungi is called Mycology.
- **2.** Fungi is chlorophyll less, central carrier tissue less, Thalophyte.
- **3.** Accumulated food in fungi remains as Glycogen.
- **4.** Its cell wall is made up of chitin. Ex. Albugo, Phytophthora Mucor etc. Fungi may creates serious diseases in plants. Most damage is caused by rust and smut. Main Fungal diseases in plants are as:

 White rust of crucifer, Loose smut of wheat, Rust of wheat, early Blight of potato, Red rot of sugarcane, Tikka diseases of ground nut, Wart disease of potato, Brown leaf spot of rice. Late blight of potato, Damping off of seedlings etc.

Bryophyta

This is the first group of land plants. In this division approximately 25000 species are included.

- 1. In byophyta there is lack of Xylem and phloem tissue.
- 2. Plant body may be of thallus like and leafy erect structure as in moss.
- 3. They lack true roots, Stem and leaves.
- 4. This community is also called Amphibian category of the plant kingdom.

The moss namely Sphagnum is capable of soaking water 18 times of its own weight. Therefore, gardeners use it to protect from drying while taking the plants from one place to another.

The Sphagnum moss is used as fuel.

The Sphagnum moss is also used as antiseptic.

Pteridophyta

The plants of this group are mostly found in wet shady places, forests and mountains.

1. The body of plants are differentiated into root, stem, and leaves. Stem remains as normal rhizome.

- 2. Reproduction occurs by spores produced inside the sporangia.
- **3.** Gametophytic phase is short lived. The diploid zygote develops into an embryo.
- **4.** Plants of this community have conducting tissues. But Xylem does not contain Vessels and Phloem does not contain companion cells.

Examples : Ferns, Azolla, Pteridium, Lycopodium etc.

II. Phanerogamus or Floral plant

The plants in this group are well developed. All the plants in this group have flowers, fruits and seeds. The plants in this group can be classified into two subgroups – Gymnosperm and Angiosperm.

(A) Gymnosperm

- **1.** These plants are in the forms of trees and bushes. Plant body are differentiated into root, stem & leaves.
- 2. Plants are woody, perennial and tall. Plant bear naked seed.
- 3. Its tap roots are well developed.
- Pollination takes place through air.

The longest plant of the Plant kingdom, Sequoia gigentia comes under it. Its height is 120 meters. This is also called Red Wood of California.

- 1. The smallest plant is Zaimia Pygmia.
- 2. Living fossils are Cycas, Ginkgo, biloba and Metasequoia.
- 3. Ginkgo biloba is also called Maiden hair tree.
- 4. Ovules and Antherzoids of Cycas is the largest in Plant kingdom.

The pollen grains of Pinus are so much in number that later it turns into Sulphur showers.

Importance of Gymnosperm

- **1. As a food –** Sago is made by extracting the juice from the stems of Cycas. Therefore, Cycas is called Sago-palm.
- **2. Wood –** The wood of Pine, Sequoia, Deodar, Spruce etc. is used for making furniture.
- **3. Vapour oil –** We get Tarpin oil from the trees of Pine, Cedrus oil from Deodar tree and Cedcast oil from Juniperous wood.
- **4. Tannin -** It is useful in tanning and making ink.
- **5. Resin** Resin is extracted from some conical plants which are used in making varnish, polish, paint etc.

(B) Angiosperm

- 1. In the plants of this sub-group seeds are found inside the fruits.
- 2. In there plants root, leaves, flowers, fruits and seeds are fully developed. In the plants of this sub-group there is seed-coat in seeds. On the basis of number of cotyledons plants are divided into two categories —

(1) Monocotyledon and (2) Dicotyledon

Monocotyledon plants: Those plants which have only one cotyledon in seed. Example:

Dicotyledons plants : Those plants which have two cotyledon in its seed are called dicotyledons.

Example :

Name of category Name of main plants

Virus

- 1. Study of virus is virology.
- **2.** Virus was discovered by Russian scientist Ivanovsky in the year 1892. (During the tests of Mosaic disease on tobacco).

- **3.** In nature, there are ultra microscopic particle known as viruses. Virus are connecting a link between living & non living.
- **4.** It has both the characters of living and non living, so it is a connecting link between living & non living.

Characters of virus

- **1.** They became active inside a living cells.
- 2. Nucleic acids replicate themselves and they reproduce rapidly.
- 3. They cause disease like bacteria & fungi.

According to parasitic nature, virus is of three types -

- 1. Plant virus RNA is present as its nucleic acid,
- 2. Animal virus DNA or sometimes RNA is found in it.
- 3. Bacteriophage— They depend only on bacteria. They kill the bacteria. DNA is found in them. Example T-2 phage.
- **4.** In man virus cause disease like mumps, chicken pox, hepatitis, palio, AIDs and Herpes.
- **5. Bacteriophages :** Bacteriophages are those virus which infect the bacteria. Example —Tobacco mosaic virus.

Note: Those viruses in which RNA substance is found as genetic material are called Retrovirus.

Bacteria

It was discovered by Antony von Lecuwenhoek of Holland in the year 1683.

- ---> Leeuwenhoek is called the father of Bacteriology. In the year 1829 Ehrenberg called it bacteria.
- ---> The year 1843-1892 Robert Koch discovered the bacteria of Tuberculosis diseases.
- ---> The year 1812-1892 Louis Pasteur discovered the vaccine of Rabies and pasteurization of milk.

- ---> On the basis of shape, bacteria is of different types :
- **1. Bacillus :** This is rod-like or cylindrical.
- 2. Round or Cocus: These are round and the smallest bacteria.
- **3. Comma shaped or Vibrio :** Like the English sign (,), example Vibrio cholerae etc.
- 4. Spirillum: Spring or screw shaped.
- ---> Some species of Azotobacter, Azospirillum and Clostridium bacteria live freely in the soil and fix atmospheric nitrogen into the nitrogenous compound. Anabaena and Nostoc cynobacteria fix atmospheric nitrogen into soil.
- ---> The species of Rhizobium and Bradyrhizobium etc. bacteria live in the roots of the Leguminous plants capable of converting atmospheric nitrogen into its compound.

Note: To preserve the milk for many days pasteurization is done. There are two methods of pasteurization —

- (a) Low temperature holding method (LTH): Milk is boiled at 62.8 degree Celsius for 30 minutes.
- **(b)** High Temperature short time method (HTSt): Milk is boiled at 71.7 degree Celsius for 15 seconds.
- ---> In leather industry separation of hair and fat from leather is done by bacteria. This is called tanning of leather.
- ---> Pickles, syrup is kept in salt or in dense liquid of sugar so that in case of bacterial attack bacteria are plasmolysed and destroyed. Therefore, pickles etc. do not get spoiled soon and can be preserved for long time.
- ---> In the Cold Storage objects are kept at low temperature (-10 degree Celsius to -18 degree Celsius).
- ---> Mycoplasma : Smallest known prokaryotic cell causing pleuropneumonia. It is also known as PPLO



JK Chrome | Employment Portal




Rated No.1 Job Application of India

Sarkari Naukri
Private Jobs
Employment News
Study Material
Notifications













JK Chrome

jk chrome Contains ads



www.jkchrome.com | Email : contact@jkchrome.com